

In-class transition (iCT) from bortezomib (V)-based induction to oral ixazomib-lenalidomide-dexamethasone vs parenteral V-based therapy: comparative effectiveness in newly diagnosed multiple myeloma



To view an electronic version of this poster, scan this QR code or visit <https://www.ctdjournal.com/2022/05/02/196>. Copies of this poster obtained through QR codes are for personal use only and may not be reproduced without permission from the congress and the author of this poster.

Caitlin L. Costello,¹ Ruemu E. Birhiray,² Suman Kambhampati,³ Joshua Richter,⁴ Rafat Abonour,⁵ Hans C. Lee,⁶ Yong Jin Kim,⁷ Kaili Ren,⁸ Dawn Marie Stull,⁹ Dasha Cherepanov,⁸ Kimberly Bogard,⁹ Stephen J. Noga,⁹ Saulius Girmius,¹⁰ Robert M. Rifkin¹¹

¹Department of Medicine, Division of Blood and Marrow Transplantation, Moores Cancer Center, University of California San Diego, La Jolla, CA, USA; ²Hematology Oncology of Indiana/American Oncology Network, Indianapolis, IN, USA; ³Kansas City Veterans Affairs Medical Center, Kansas City, MO, USA; ⁴Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA; ⁵Indiana University School of Medicine, Indianapolis, IN, USA; ⁶M.D. Anderson Cancer Center, Houston, TX, USA; ⁷Evidera, Data Analytics, St-Laurent, Quebec, Canada; ⁸Takeda Development Center Americas, Inc. (TDCA), Lexington, MA, USA; ⁹Takeda Pharmaceuticals U.S.A., Inc., Lexington, MA, USA; ¹⁰TriHealth Cancer Institute, Cincinnati, OH, USA; ¹¹Rocky Mountain Cancer Centers/US Oncology Research, Denver, CO, USA

Background

- Long-term proteasome inhibitor (PI)-based treatment can improve outcomes for patients with multiple myeloma (MM)^{1,2}
- However, prolonged parenteral PI therapy (e.g., with bortezomib [V]) can be challenging to achieve in routine clinical practice, and outcomes for patients are often poorer in this setting compared with clinical trials³
 - This discrepancy could be due in part to the fact that up to 72% of real-world patients with relapsed/refractory MM (RRMM) would not meet the eligibility criteria for randomized clinical trials⁴
 - Real-world studies with less stringent eligibility criteria may permit inclusion of a more diverse patient population and better inform on the effectiveness of therapies as used in routine clinical practice
 - Various physical, geographical, and/or socioeconomic barriers to prolonged therapy with parenteral PIs are encountered in community practice,³ including:
 - Burden of repeated intravenous/subcutaneous administration
 - Difficulty traveling to treatment centers
 - Patient preference for treatment outside of a clinic
 - Comorbidities
 - Toxicities
- The phase 4, community-based, single-arm US MM-6 study (NCT03173092) is assessing in-class transition (iCT) from V-based induction to all-oral ixazomib-lenalidomide-dexamethasone (IRd) in transplant ineligible newly-diagnosed MM (NDMM) patients treated in routine clinical practice, with the objective of increasing the duration of PI-based treatment while maintaining quality of life⁵
 - The US MM-6 study provided the patients for the IRd cohort
- INSIGHT MM is the largest global, prospective, observational study (NCT02761187) of MM patients (>4200),⁶ and provided a subset of patients on V-based therapy as the comparator cohort
- This enabled assessment of iCT versus V-based therapy in NDMM patients in community oncology practices in the US


Aims

- To examine the comparative effectiveness of IRd following initial V-based induction (3 cycles; US MM-6 patients; 'IRd cohort') versus patients who continued to receive V-based therapy (INSIGHT MM patients; 'V-based cohort') in patients with NDMM

Methods

- Patients**
- Transplant ineligible US NDMM patients with \geq stable disease (SD) after 3 cycles of V-based induction and baseline Eastern Cooperative Oncology Group performance status (ECOG PS) of 0, 1 or 2 from the US MM-6 and INSIGHT MM studies^{5,6}
- Endpoints**
- Study outcomes included first-line duration of treatment (DOT), overall response rate (ORR), progression-free survival (PFS), overall survival (OS), and reasons for treatment discontinuation
- Statistical Analyses**
- These secondary analyses were weighted using the inverse probability of treatment weighting (IPTW) approach to reduce the imbalance of potential confounding factors between the two cohorts (Figure 2)
 - The Kaplan-Meier method was used to examine DOT, PFS, OS, and associated 95% confidence intervals (CIs); the log-rank test was used to compare distribution of time to events
 - The Clopper-Pearson method was applied to estimate 95% CIs for ORR
 - Statistical significance was evaluated at $\alpha=0.05$


Figure 2: Populations for the IRd and V-based cohorts selected from US MM-6 and INSIGHT MM




¹ECOG PS was only measured at baseline and then on a yearly basis in INSIGHT MM. ²Frailty status was assessed based on the Simplified Frailty Score estimated using age, CCI and ECOG PS. ³CCI, Charlson Comorbidity Index; ISS, International System Staging; NCCN, National Comprehensive Cancer Network; PD, progressive disease; VcD, bortezomib-cyclophosphamide-dexamethasone; VRcD, bortezomib-lenalidomide-cyclophosphamide-dexamethasone; VRd, bortezomib-lenalidomide-dexamethasone

Question

In transplant ineligible patients with NDMM, does IRd following 3 initial cycles of V-based induction (US MM-6 patients; 'IRd cohort') have comparable effectiveness with patients who continued to receive V-based therapy (INSIGHT MM patient; 'V-based cohort')?



Results



*IPTW-weighted cohorts. [†]Defined as the proportion of patients with partial response (PR), very good partial response (VGPR), complete response (CR), and stringent complete response (sCR) during initial treatment regimen and prior to disease progression. [‡]Time from the index date (date that patients began V-based therapy) to the date of the last administration of any of the three study drugs in the IRd regimen (IRd cohort) or first-line V-based regimen (V-based cohort) (event), death (due to any cause, event), or end of follow-up (censored).

Key take aways

US MM-6 NDMM patients who transitioned to IRd after 3 initial cycles of V-based induction had a significantly higher ORR and longer DOT compared with patients who continued to receive V-based therapy in INSIGHT MM. The results suggest that iCT from parenteral V-based therapy to all-oral IRd may improve outcomes in patients treated at community oncology practices.

Results

- Patients**
- 100 patients from the IRd cohort (US MM-6) and 111 patients from the V-based cohort (INSIGHT MM) were included in the analysis
 - In the IPTW-weighted cohorts, 113 patients from MM-6 and 102 patients from INSIGHT MM contributed to the analyses
 - In the weighted IRd versus V-based cohorts: median age was 75.0 years versus 74.8 years; 57% versus 51% of patients were male; 37% versus 29% had an ECOG PS of \leq 2; 49% versus 41% had ISS stage III at initial diagnosis, and 79/18/3% versus 77/20/3% of patients had received VRd/VcD/VRcD as initial induction therapy (Table 1)
- Efficacy**
- Adjusted ORRs in the IRd versus V-based cohorts were 73.2% (95% CI 65.0–81.3) versus 57.5% (95% CI 47.9–67.1; p<0.0001) (Figure 1A)
 - After a median follow-up of 20.3 and 15.8 months in the IRd and V-based cohorts, respectively, median DOT was 10.8 months (95% CI 6.5–24.4) versus 5.3 months (95% CI 4.3–7.0; p<0.0001) (Figure 1B)
 - Median PFS and OS were not estimable (NE) in either cohort (Figures 3 & 4)
 - PFS rates (95% CI) for the IRd vs V-based cohorts were:
 - 6-month: 96.0% (81.2–99.2) versus 95.3% (87.1–98.4)
 - 12-month: 86.9% (69.6–94.7) versus 87.5% (76.9–93.5)
 - 18-month: 85.7% (68.1–94.0) versus 80.1% (67.8–88.1)
 - 24-month: 85.7% (68.1–94.0) versus 76.5% (62.6–85.8)
 - OS rates (95% CI) for the IRd vs V-based cohorts were:
 - 6-month: 100% (100–100) versus 98.0% (90.5–99.6)
 - 12-month: 99.3% (61.0–100) versus 96.6% (88.2–99.1)
 - 18-month: 96.9% (80.9–99.5) versus 90.4% (79.7–95.6)
 - 24-month: 94.0% (77.7–98.5) versus 84.9% (70.6–92.6)

Reasons for Treatment Discontinuation

- In the IRd and V-based cohorts, 18% and 24% of patients discontinued IRd and V, respectively, due to an adverse event (AE; Table 2)

Figure 3: Adjusted PFS in the IRd and V-based cohorts




Figure 4: Adjusted OS in the IRd and V-based cohorts




Table 2: Reasons for discontinuation of IRd or V

	IRd cohort (N=113*)	V-based cohort (N=102*)
Discontinued medication, n (%)	72 (63.3)	73 (71.9)
Reason for discontinuation, n (%)		
PD	16 (22.6)	9 (12.8)
AE	13 (17.6)	18 (24.4)
Patient request	34 (48.0)	1 (0.8)
Planned therapy completed	-	33 (44.6)
Inadequate response	-	1 (1.8)
Other	9 (11.9)	11 (15.5)

Limitations

- This study has limitations, of which some are inherent to analyses of observational data
- INSIGHT-MM (V-based cohort) was an observational study where outcomes were assessed on a quarterly basis, whereas outcomes for the US MM-6 single arm clinical trial (IRd cohort) were assessed at the end of each 28-day treatment cycle
 - This difference in timing of the assessments may have introduced bias, as patients in the IRd cohort were being followed-up more frequently
- Across treatment cohorts there were differences in geography (INSIGHT-MM was a global study, whereas US MM-6 was restricted to the US) and type of treatment centers (most treatment centers in INSIGHT were academic, whereas US MM-6 centers were from community settings)
 - To account for the differences in geography, only INSIGHT-MM patients residing in the US were included in this analysis. This ensured more balanced cohorts, however resulted in a reduced sample size for this analysis
- Propensity score values used in the IPTW were based on a vector of observed covariates, and thus any confounding due to unobserved covariates (e.g., residual confounding) could not be accounted for

Conclusions

- US MM-6 NDMM patients who transitioned to IRd after 3 initial cycles of V-based induction had a significantly higher ORR and longer DOT compared with patients who continued to receive V-based therapy in INSIGHT MM
- iCT from V-based therapy to all-oral IRd therapy resulted in a lower treatment discontinuation rate due to AEs compared with patients who continued to receive V-based therapy
- The results suggest that iCT from V-based therapy to all-oral IRd may improve outcomes in patients treated at community oncology clinics
- In addition to offering a viable treatment option, the ability to transition from a parenteral to an oral treatment regimen could prevent disruption to patients' treatment course
- All-oral IRd may be most beneficial to patients with restricted mobility (e.g. elderly patients), those who may prefer to remain outside of a hospital/clinic setting for treatments, or those with work commitments/family obligations

References

- Mateos MV, et al. *Am J Hematol* 2015;90:314–9
- Jimenez-Zepeda VH, et al. *Ann Hematol* 2017;96:431–9
- Richardson PG, et al. *Blood Cancer J* 2018;8:109
- Charl A, et al. *Clin Lymphoma Myeloma Leuk* 2020;20:8–17
- Manda S, et al. *Clin Lymphoma Myeloma Leuk* 2020;20:e910–25
- Costello C, et al. *Future Oncol* 2019;15:1411–28
- Facon T, et al. *Leukemia* 2020;34:224–33
- Thomas C & Thomas L. *Dtsch Arztebl Int* 2009;106:849–54

Acknowledgments

These studies were funded by Takeda Development Center Americas, Inc. (TDCA), Lexington, MA, USA. The authors would like to thank all patients and their families, as well as investigators at all clinical sites for their participation in the study. Medical writing support for the development of this poster, under the direction of the authors, was provided by Laura Webb, PhD, of Ashfield MedComms, an Inizio Company, funded by Takeda Pharmaceuticals U.S.A., Inc., Lexington, MA and complied with the Good Publication Practice-3 (GPP3) guidelines (Battisti WP, et al. *Ann Intern Med* 2015;163:461–4). We also thank Michael Stokes, of Evidera, Data Analytics, for his contribution towards study design and data analysis.

Disclosures

CLC: Member of advisory council/committees for BMS, Janssen, Takeda, and Pfizer; REB: Member of advisory council/committees for Array Biopharma Inc., Lilly Oncology, Janssen Scientific Affairs Lic, Epizyme, Tg Therapeutics, and Regeneron; Other financial relationships with Janssen Biotech Inc., Amgen Inc., Puma Biotechnology Inc., Lilly USA Lic, Incyte Corporation, Pharmacyclics Lic (an Abbvie company), Genzyme Corporation, Dova/Sobi Pharmaceuticals, Exelixis Inc., E.R. Squibb & Sons Lic, Abbvie Inc., AstraZeneca Pharmaceuticals Lp, Sanofi, Daiichi Sancho, Morphosys, Regeneron, Glaxo Oncology, and Seagen; SK: none; JR: Consulting fees from Takeda, and BMS/Celgene; RA: Member of advisory council/committees for Takeda, BMS, and Jensen; Honoraria from Takeda, BMS, Jensen, and GSK; Consulting fees from Takeda, BMS, Jensen, and GSK; Grants or funds from Takeda, BMS, Jensen, and GSK;

HCL: Consulting fees from BMS/Celgene, Genentech, Janssen, Karyopharm, Legend Biotech, GSK, Sanofi, Pfizer, Monte Rosa Therapeutics, Immunitis Therapeutics, Oncopetides, and Takeda Pharmaceuticals; Grants or funds from Amgen, Janssen, GSK, Regeneron, and Takeda Pharmaceuticals; YJK: Employment with Takeda Pharmaceutical Company Limited; Consulting fees from Takeda; KR: Employment with Takeda Development Center Americas, Inc. (TDCA); DMS: Employment with Takeda Pharmaceuticals; Ownership of stock/shares with Takeda Pharmaceuticals; DC: Employment with Takeda; Ownership of stock/shares with Takeda; KB: Employment with Takeda Oncology; SJN: Employment with Takeda Pharmaceuticals; SG: Member of advisory council/committees for Takeda, BMS, Janssen, and GSK; Honoraria from Takeda, BMS, Beigene, GSK, and Adaptive; RMR: Employment with McKesson; Ownership of stock/shares with McKesson; Member of advisory council/committees for Amgen, BMS, Coherus, Fresenius, and Takeda.