



# Real-World Treatment Patterns and Effectiveness in Patients with ALK+ Advanced NSCLC Treated With 1L ALK TKIs

Yuanbin Chen<sup>1</sup>, Huamao Mark Lin<sup>2</sup>, Yin Wan<sup>2</sup>, Yu Yin<sup>2</sup>, Yanyu Wu<sup>2</sup>, Luis Hernandez<sup>3</sup>, Christopher Danes<sup>3</sup>, Eric N. Churchill<sup>3</sup>, Anastasios Dimou<sup>4</sup>

<sup>1</sup>Cancer & Hematology Centers of Western Michigan, Grand Rapids, MI, USA; <sup>2</sup>Takeda Development Center Americas, Inc., Lexington, MA, USA; <sup>3</sup>Takeda Pharmaceuticals U.S.A. Inc., Lexington, MA, USA; <sup>4</sup>Mayo Clinic College of Medicine, Rochester, MN, USA

## Background

- Anaplastic lymphoma kinase-positive (ALK+) non-small cell lung cancer (NSCLC) accounts for approximately 4–5% of all NSCLC cases.<sup>1,2</sup>
- Approved targeted therapies for ALK+ NSCLC include the first-generation ALK tyrosine kinase inhibitor (TKI), crizotinib, and next-generation ALK TKIs, such as brigatinib, alectinib, and lorlatinib.<sup>3-6</sup>
  - For first-line (1L) treatment of advanced ALK+ NSCLC, crizotinib, brigatinib, alectinib, and lorlatinib received Food and Drug Administration (FDA) approvals in January 2013, May 2020, November 2017, and March 2021, respectively.<sup>4</sup>
- In their respective phase III clinical trials, brigatinib, alectinib, and lorlatinib exhibited superior clinical efficacy compared with crizotinib for the 1L treatment of patients with ALK+ NSCLC.<sup>8-11</sup>
- There is a lack of real-world data on treatment patterns and clinical outcomes in the 1L setting for patients with ALK+ advanced NSCLC.

## Objective


- To describe real-world treatment patterns and effectiveness in patients with ALK+ advanced NSCLC treated with 1L ALK TKIs.

## Methods

- Study design and patients**
- This retrospective cohort study used data extracted from the US Flatiron electronic medical record-derived database (Jan 2011–Sept 2023) (Figure 1).
  - Patients were included if they:
    - Had a diagnosis of ALK+ advanced NSCLC;
    - Received alectinib, brigatinib, crizotinib, or lorlatinib monotherapy as their first ALK TKI treatment after initial advanced NSCLC diagnosis; and
    - Were aged ≥18 years at the time of initiating 1L alectinib, brigatinib, crizotinib, or lorlatinib monotherapy.
  - Patients with a positive ROS1 test result anytime were excluded.
  - Index date was defined as the date of initiation of 1L ALK TKI.
- Study outcomes and statistical analyses**
- Study outcomes included first subsequent treatment (second-line, 2L) after 1L ALK TKI; real-world time to treatment discontinuation (rwTTD); and real-world time to next treatment (rwTNT).
  - Kaplan-Meier survival analyses were used to estimate the median rwTTD and rwTNT for each study cohort.
  - A multivariate Cox proportional hazard model was performed to compare rwTTD and rwTNT for 1L brigatinib or 1L alectinib vs 1L crizotinib, adjusting for relevant covariates (age, sex, race, Eastern Cooperative Oncology Group performance score [ECOG PS], smoking, baseline brain metastasis, time from advanced diagnosis to index date).
  - To further compare the real-world effectiveness of 1L brigatinib vs 1L alectinib or 1L crizotinib, propensity score matching using an inverse probability of treatment weighting (IPTW) method was applied; adjusted covariates used for propensity matching included age, sex, smoking status, ECOG PS, baseline brain metastasis, and time from advanced diagnosis to index date.
  - A sensitivity analysis was conducted to minimize bias resulting from different FDA approval dates of the ALK TKIs, which included patients treated with alectinib, brigatinib, or crizotinib on or after May 22, 2020 (the FDA approval date for brigatinib).
  - Formal assessments on the outcomes (TTD/TTNT) of 1L lorlatinib were not conducted due to the small sample size and immature data.

## Results

- Patients**
- The study included 832 patients, of which 643 patients received 1L alectinib, 28 received 1L brigatinib, 144 received 1L crizotinib, and 17 received 1L lorlatinib (Figures 1 and 2).
  - Mean (standard deviation [SD]) age of patients in the 1L alectinib, 1L brigatinib, 1L crizotinib, and 1L lorlatinib cohorts were 61.8 (13.2), 61.1 (14.1), 70.0 (11.5), and 60.6 (11.3) years, and 44.0%, 57.1%, 45.1%, and 35.3% of patients were male, respectively (Table 1).



1L, first-line; ALK, anaplastic lymphoma kinase; NSCLC, non-small cell lung cancer; TKI, tyrosine kinase inhibitor.

### Question

What are the real-world treatment patterns and effectiveness in patients with ALK+ advanced NSCLC treated with 1L ALK TKIs?

---

### Study Design

**Figure 1. Study design**

1L, first-line; ALK, anaplastic lymphoma kinase; RWD, real-world data; rwTTD, real-world time to treatment discontinuation; rwTNT, time to next treatment; TKI, tyrosine kinase inhibitor; U.S., United States.

1L Brigatinib (n=28)  
1L Alectinib (n=643)  
1L Crizotinib (n=144)  
1L Lorlatinib (n=17)

**Outcomes:** treatment patterns, rwTTD, and rwTNT

---

### Results

**Figure 3. Kaplan-Meier curves for adjusted rwTTD for (A) 1L alectinib vs. 1L brigatinib, and (B) 1L brigatinib vs. 1L crizotinib, and adjusted rwTNT for (C) 1L alectinib vs. 1L brigatinib, and (D) 1L brigatinib vs. 1L crizotinib**

1L, first-line; DOT, duration of treatment; rwTTD, real-world time to treatment discontinuation; rwTNT, time to next treatment.

### Key Takeaways

- This real-world study showed that treatment with 1L alectinib and 1L brigatinib had similar rwTTD and rwTNT.
- The real-world findings from this study mirror and support the phase III ALTA-1L trial data demonstrating the efficacy of 1L brigatinib vs. crizotinib in patients with ALK+ advanced NSCLC.

**Table 1. Baseline characteristics**

Baseline characteristics	1L alectinib n=643	1L brigatinib n=28	1L crizotinib n=144	1L lorlatinib n=17
Age at index date, years, mean (SD)	61.6 (13.2)	61.1 (14.1)	70.0 (11.5)	60.6 (11.3)
Male, n (%)	283 (44.0)	16 (57.1)	65 (45.1)	6 (35.3)
Race, n (%)				
Asian	48 (7.5)	2 (7.1)	8 (5.6)	0
Black or African American	47 (7.3)	1 (3.6)	14 (9.7)	4 (23.5)
Hispanic or Latino	1 (0.2)	0	0	0
Other/Unknown	143 (22.2)	14 (50.0)	25 (17.4)	5 (29.4)
White	404 (62.8)	11 (39.3)	97 (67.4)	8 (47.1)
History of smoking, n (%)	273 (42.5)	8 (28.6)	100 (69.4)	4 (23.5)
Patients with brain metastasis at baseline, n (%)	97 (15.1)	5 (17.9)	29 (20.1)	6 (35.3)
ECOG PS at index date, n (%)				
0	215 (33.4)	8 (28.6)	28 (19.4)	11 (64.7)
1	228 (35.5)	6 (21.4)	59 (41.0)	0
2	60 (9.3)	1 (3.6)	21 (14.6)	0
3	13 (2.0)	0	8 (5.6)	1 (5.9)
4	2 (0.3)	1 (3.6)	1 (0.7)	0
Unknown	125 (19.4)	12 (42.9)	27 (18.8)	5 (29.4)
Time from initial advanced diagnosis to 1L ALK TKI, months, median (IQR)	1.5 (0.9, 4.4)	2.5 (1.2, 44.6)	7.5 (2.4, 18.7)	4.0 (1.6, 35.6)
Prior non-ALK TKI treatments after initial advanced diagnosis, N (%)				
Chemo	33 (5.1)	1 (3.6)	5 (3.5)	1 (5.9)
Chemo+others	5 (0.8)	0	5 (3.5)	0
IO	8 (1.2)	0	8 (5.6)	0
IO+chemo/other agents	79 (12.3)	3 (10.7)	52 (36.1)	0
Others	9 (1.4)	0	11 (7.6)	2 (11.8)
No prior non-ALK TKI therapy, n (%)	509 (79.2)	24 (85.7)	63 (43.8)	14 (82.4)
Number of prior lines of therapy <sup>a</sup> (%)				
1	94 (14.6)	2 (7.1)	40 (27.8)	3 (17.6)
2	27 (4.2)	1 (3.6)	20 (13.9)	0
≥3	17 (2.6)	1 (3.6)	22 (15.3)	0
Follow-up duration from start of 1L ALK TKI treatment until death or last activity date, months				
Mean (SD)	24.4 (19.3)	17.1 (10.8)	14.1 (15.2)	13.7 (8.1)

<sup>a</sup>All prior therapies were non-ALK TKI drugs.

1L, first-line; ALK, anaplastic lymphoma kinase; chemo, chemotherapy; ECOG PS, Eastern Cooperative Oncology Group performance score; IO, immunology; IQR, interquartile range; SD, standard deviation; TKI, tyrosine kinase inhibitor.

**REFERENCES**

1. Barlesi F, et al. *Lancet*. 2016;387(10026):1415-1426.
2. Solomon B, et al. *J Thorac Oncol*. 2009; 4(12):1450-1454.
3. Spagnuolo A, et al. *Expert Opin Emerg Drugs*. 2018;23(3):231-241.
4. Peng L, et al. *Front Oncol*. 2022;12:863461.
5. XALKOR® (crizotinib) [prescribing information]. Pfizer, New York, NY, 2023.
6. ALUNBRIG® (brigatinib) [prescribing information]. Takeda Pharmaceuticals America, Inc., Lexington, MA, 2022.
7. ALECENSA® (alectinib) [prescribing information]. Genentech USA, Inc., South San Francisco, CA, 2024.
8. LORBRENA® (lorlatinib) [prescribing information]. Pfizer, New York, NY, 2023.
9. Camidge DR, et al. *J Thorac Oncol*. 2021;16(12):2091-2108.
10. Peters S, et al. *New Engl J Med*. 2017;377(9):829-838.
11. Solomon BJ, et al. *J Clin Oncol*. 2024;JCO2400581.

**Table 2. Distribution of first subsequent treatment following 1L treatments**

n (%) patients with subsequent treatment	1L alectinib n=643	1L brigatinib n=28	1L crizotinib n=144	1L lorlatinib n=17
None	410 (63.8)	21 (75.0)	82 (56.9)	14 (82.4)
Chemo	10 (1.6)	0	16 (11.1)	0
Chemo+IO	8 (1.2)	0	6 (4.2)	0
IO	8 (1.2)	0	8 (5.6)	0
TKI	186 (28.9)	6 (21.4)	13 (9.0)	2 (11.8)
ALK TKI-based regimen <sup>a</sup>	32 (5.0)	0	5 (3.5)	1 (5.9)
Alectinib monotherapy	8 (1.2)	1 (3.6)	6 (4.2)	0
Brigatinib monotherapy	36 (5.6)	0	1 (0.7)	1 (5.9)
Ceritinib monotherapy	4 (0.6)	0	0	0
Crizotinib monotherapy	2 (0.3)	0	2 (1.4)	0
Lorlatinib monotherapy	112 (17.4)	5 (17.9)	0	0
Others	13 (2.0)	1 (3.6)	18 (12.5)	1 (5.9)

<sup>a</sup>Refers to ALK TKI in combination with other agents such as chemo and IO.


1L, first-line; ALK, anaplastic lymphoma kinase; chemo, chemotherapy; IO, immunology; TKI, tyrosine kinase inhibitor.

**Table 3. Unadjusted and adjusted rwTTD and rwTNT**

1L ALK TKI <sup>a</sup>	N	Unadjusted median <sup>b</sup> months (95% CI)	Adjusted median <sup>b</sup> months (95% CI)	Adjusted HR <sup>c</sup> (95% CI)	P value
<b>rwTTD</b>					
1L alectinib	643	23.2 (18.2–29.2)	23.5 (18.6–34.2)	0.304 (0.241–0.384)	<0.0001
1L brigatinib	28	23.9 (8.4–NR)	23.9 (8.4–NR)	0.357 (0.199–0.641)	0.0006
1L crizotinib	144	4.1 (3.0–5.0)	6.7 (3.0–15.7)	-	-
<b>rwTNT</b>					
1L alectinib	643	25.5 (20.5–31.6)	28.9 (22.8–36.9)	0.321 (0.254–0.405)	<0.0001
1L brigatinib	28	23.9 (10.8–NR)	23.9 (10.8–NR)	0.377 (0.210–0.675)	0.001
1L crizotinib	144	5.3 (4.1–7.4)	12.5 (NR–NR)	-	-

<sup>a</sup>1L lorlatinib was not assessed due to small sample size. <sup>b</sup>Estimated using Kaplan-Meier survival analysis. <sup>c</sup>Post-IPTW analysis for brigatinib vs. alectinib and brigatinib vs. crizotinib. <sup>d</sup>Cox proportional hazard model with crizotinib as control.

1L, first-line; ALK, anaplastic lymphoma kinase; CI, confidence interval; HR, hazard ratio; IPTW, inverse probability of treatment weighting; NR, not reached; rwTTD, real-world time to treatment discontinuation; rwTNT, time to next treatment; TKI, tyrosine kinase inhibitor.



<sup>a</sup>Included patients treated with alectinib, brigatinib, or crizotinib on or after May 22, 2020 (the FDA approval date for brigatinib). DOT, duration of therapy; FDA, Food and Drug Administration; rwTTD, real-world time to treatment discontinuation; rwTNT, time to next treatment.

## Limitations

- The sample size for the 1L brigatinib cohort is small; larger sample size and longer follow-up are warranted for future studies.
- Data generated from real-world clinical practice are subject to miscoding, errors, underreporting, or missing values.
  - ECOG data were missing across the treatment cohorts, with a high percentage of missing data for the 1L brigatinib cohort.
  - Information on interventions that happened outside of the Flatiron network were not captured.
  - The majority of the clinics or institutions within the Flatiron network are community-based practices; therefore, the results may not be generalizable to academic institutions.

## Discussion

- In patients with ALK+ advanced NSCLC, treatment with 1L alectinib and 1L brigatinib was found to have similar rwTTD and rwTNT.
- 1L alectinib and 1L brigatinib were associated with improved rwTTD and rwTNT vs 1L crizotinib.
- IPTW analysis showed similar effectiveness for 1L alectinib and 1L brigatinib and improved effectiveness for 1L brigatinib and 1L alectinib vs 1L crizotinib.

**DISCLOSURES**  
Yin Wan reports employment with Takeda Development Center Americas, Inc. (TDCA).  
**ACKNOWLEDGEMENTS**  
This study was funded by Takeda Development Center Americas, Inc. (TDCA). Medical writing and editorial support was provided by SNELL Medical Communication, Inc. and funded by TDCA.